



Complete Genome Sequence of *Nitrosomonas cryotolerans* ATCC 49181, a Phylogenetically Distinct Ammonia-Oxidizing Bacterium Isolated from Arctic Waters

Marlen C. Rice,^a Jeanette M. Norton,^a Lisa Y. Stein,^b Jessica Kozlowski,^b Annette Bollmann,^c Martin G. Klotz,^{d,e} Luis Sayavedra-Soto,^f Nicole Shapiro,^g Lynne A. Goodwin,^h Marcel Huntemann,^g Alicia Clum,^g Manoj Pillay,^g Neha Varghese,^g Natalia Mikhailova,^g Krishna Palaniappan,^g Natalia Ivanova,^g Supratim Mukherjee,^g T. B. K. Reddy,^g Chew Yee Ngan,^g Chris Daum,^g Nikos Kyrpides,^g Tanja Woyke^g

Utah State University, Logan, Utah, USA^a; University of Alberta, Edmonton, Alberta, Canada^b; Miami University, Oxford, Ohio, USA^c; Queens College in The City University of New York, Flushing, New York, USA^d; The Institute of Marine Microbes and Ecospheres, Xiamen University, Xiamen, China^e; Oregon State University, Corvallis, Oregon, USA^f; DOE Joint Genome Institute, Walnut Creek, California, USA^g; Los Alamos National Laboratory, Bioscience Division, Los Alamos, New Mexico, USA^h

ABSTRACT *Nitrosomonas cryotolerans* ATCC 49181 is a cold-tolerant marine ammonia-oxidizing bacterium isolated from seawater collected in the Gulf of Alaska. The high-quality complete genome contains a 2.87-Mbp chromosome and a 56.6-kbp plasmid. Chemolithoautotrophic modules encoding ammonia oxidation and CO₂ fixation were identified.

Nitrosomonas cryotolerans ATCC 49181 was isolated from surface seawater at 59°47'67"N, 151°55'08"W in 1980 and described in 1988 (1). *Nitrosomonas cryotolerans* is an obligate halophilic betaproteobacterium in the *Nitrosomonadaceae* (2) that is able to grow at temperatures as low as -5°C (3), the coldest temperature known for any bacterium in the *Nitrosomonadaceae*; *Nitrosomonas cryotolerans* was selected for sequencing based on phylogenetic and physiologic interests. Genome comparisons with *N. cryotolerans* will lead to better delineation of the *Nitrosomonas* genus.

Nitrosomonas cryotolerans was grown in marine medium containing 25 mM ammonium at 15°C (4). Cultures were harvested by centrifugation, and genomic DNA (gDNA) was isolated using the JGI cetyltrimethylammonium bromide (CTAB) protocol. The quality of gDNA was checked by gel electrophoresis and by amplification and sequencing of the V4 region of the small subunit (SSU) rRNA gene (5).

Genomic DNA was sequenced at the Department of Energy (DOE) Joint Genome Institute (JGI) using Pacific Biosciences (PacBio) technology (6). A PacBio SMRTbell library was constructed and sequenced on the PacBio RS platform, which generated 178,102 reads totaling 560.1 Mb. The input read coverage was 191.1×. Raw reads were assembled using HGAP version 2.2.0.p1 (7). The final assembly contains two contigs (one chromosome and one plasmid) totaling 2.87 Mbp in size. An earlier version accomplished using the Illumina HiSeq 2000 platform remained in 91 scaffolds, with nearly identical repeats of key gene clusters unresolved, and was not combined with the PacBio sequence for the complete genome reported here.

Genes were identified using Prodigal (8), using the DOE-JGI microbial annotation pipeline (9), followed by manual curation using GenePRIMP (10). The predicted coding sequences (CDSs) were translated and used to search the NCBI nonredundant, UniProt,

Received 10 January 2017 Accepted 12 January 2017 Published 16 March 2017

Citation Rice MC, Norton JM, Stein LY, Kozlowski J, Bollmann A, Klotz MG, Sayavedra-Soto L, Shapiro N, Goodwin LA, Huntemann M, Clum A, Pillay M, Varghese N, Mikhailova N, Palaniappan K, Ivanova N, Mukherjee S, Reddy TBK, Yee Ngan C, Daum C, Kyrpides N, Woyke T. 2017. Complete genome sequence of *Nitrosomonas cryotolerans* ATCC 49181, a phylogenetically distinct ammonia-oxidizing bacterium isolated from Arctic waters. Genome Announc 5:e00011-17. <https://doi.org/10.1128/genomeA.00011-17>.

Copyright © 2017 Rice et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Jeanette M. Norton, jeanette.norton@usu.edu.

TIGRFam, Pfam, KEGG, COG, and InterPro databases. Transfer RNA genes were identified using the tRNAscan-SE tool (11). Ribosomal RNA genes were found by searches against models from SILVA (12). Other noncoding RNAs were found using INFERNAL (13). Further gene prediction and manual curation were performed within the JGI Integrated Microbial Genomes platform (14, 15).

The genome of *N. cryotolerans* contains one ribosomal operon (SAMN02743940_1059 to SAMN02743940_1062). Metabolic modules encoding chemolithotrophic ammonia catabolism include three clusters each of ammonia-monooxygenase and hydroxylamine-oxidoreductase genes. The *amo* gene clusters have unique arrangements of neighboring genes, including the *amoCABEDcopCD* cluster (SAMN02743940_0955 to SAMN02743940_0961), *amoCABED* cluster (SAMN02743940_1533 to SAMN02743940_1537), and *amoCAB* cluster (SAMN02743940_2438 to SAMN02743940_2436). The genome contains 2 standalone *amoC* genes (SAMN02743940_2312 and 2350). Three *haoABcycAB* clusters are located at SAMN02743940_0346 to SAMN02743940_0349, SAMN02743940_0686 to SAMN02743940_0689, and SAMN02743940_1793 to SAMN02743940_1796. The *N. cryotolerans* genome contains a single gene cluster encoding the Calvin-Benson-Bassham cycle for carbon assimilation (SAMN02743940_1712 to SAMN02743940_1708 *LysR*, *CbbL*, *CbbS*, *CbbQ*, and *CbbO*) with highest nucleic acid sequence identity to homologous genes in *Nitrosomonas ureae* Nm10 (16) and *Nitrosomonas* sp. AL212 (17) related to the form 1A (green-like) subgroup. *Nitrosomonas cryotolerans* contains a cluster encoding urease (SAMN02743940_1408 to SAMN02743940_1413) and a urea transporter (SAMN02743940_1414). We identified an *nirK* gene at SAMN02743940_0821, putatively involved in nitrite processing, and a nitric oxide reductase *norCBQD* cluster at SAMN02743940_1672 to SAMN02743940_1675 suggestive of nitrifier denitrification. The nitrosocyanin protein, unique to ammonia-oxidizing bacteria (AOB), was encoded by SAMN02743940_2670. The evolutionary relationships in the *Nitrosomonadaceae* are currently under reconsideration.

Accession number(s). This complete genome sequence is deposited in ENA under accession numbers [FSRO01000001](#) to [FSRO01000002](#).

ACKNOWLEDGMENTS

We thank JGI for resequencing this genome using improved sequencing technologies.

The work conducted was supported by the Joint Genome Institute, Office of Science of the U.S. Department of Energy under contract no. DE-AC02-05CH11231 for CSP 2010 project 1012224, by NIFA Award 2011-67019-30178, and by the Utah Agricultural Experiment Station, Utah State University and approved as journal paper UAES 8953.

J.M.N. was the principal investigator (PI) of the original CSP and lab director; sequencing, assembly, and annotation were performed at JGI, and the project manager at JGI was N.S. M.C.R. assisted in culture maintenance and wrote the first draft of the article; additional coauthors discussed, edited, and improved the proposal and manuscript.

REFERENCES

1. Jones RD, Morita RY, Koops HP, Watson SW. 1988. A new marine ammonium-oxidizing bacterium, *Nitrosomonas cryotolerans* sp.-nov. *Can J Microbiol* 34:1122–1128. <https://doi.org/10.1139/m88-198>.
2. Prosser JI, Head IM, Stein L. 2014. The family *Nitrosomonadaceae*, p 901–918. In Rosenberg E, DeLong E, Lory S, Stackebrandt E, Thompson F (ed), *The prokaryotes*. Springer-Verlag, Berlin, Heidelberg, Germany.
3. Jones RD, Morita RY. 1985. Low-temperature growth and whole-cell kinetics of a marine ammonium oxidizer. *Mar Ecol Prog Ser* 21:239–243.
4. Norton JM, Alzerreca JJ, Suwa Y, Klotz MG. 2002. Diversity of ammonia monooxygenase operon in autotrophic ammonia-oxidizing bacteria. *Arch Microbiol* 177:139–149. <https://doi.org/10.1007/s00203-001-0369-z>.
5. He Z, Xu M, Deng Y, Kang S, Kellogg L, Wu L, Van Nostrand JD, Hobbie SE, Reich PB, Zhou J. 2010. Metagenomic analysis reveals a marked divergence in the structure of belowground microbial communities at elevated CO₂. *Ecol Lett* 13:564–575. <https://doi.org/10.1111/j.1461-0248.2010.01453.x>.
6. Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, Bibillo A, Bjornson K, Chaudhuri B, Christians F, Cicero R, Clark S, Dalal R, Dewinter A, Dixon J, Foquet M, Gaertner A, Hardenbol P, Heiner C, Hester K, Holden D, Kearns G, Kong XX, Kuse R, Lacroix Y, Lin S, Lundquist P, Ma CC, Marks P, Maxham M, Murphy D, Park I, Pham T, Phillips M, Roy J, Sebra R, Shen G, Sorenson J, Tomaney A, Travers K, Trulson M, Vieceli J, Wegener J, Wu D, Yang A, Zaccarin D, et al. 2009. Real-time DNA sequencing from single polymerase molecules. *Science* 323:133–138. <https://doi.org/10.1126/science.1162986>.
7. Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Non-hybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
8. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site

- identification. *BMC Bioinformatics* 11:119. <https://doi.org/10.1186/1471-2105-11-119>.
9. Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Palaniappan K, Szeto E, Pillay M, Chen I-M, Pati A, Nielsen T, Markowitz VM, Kyrpides NC. 2015. The standard operating procedure of the DOE-JGI microbial genome annotation pipeline (MGAP v. 4). *Stand Genomic Sci* 10:86.
 10. Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. 2010. GenePRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods* 7:455–457. <https://doi.org/10.1038/nmeth.1457>.
 11. Lowe TM, Eddy SR. 1997. TRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25:955–964. <https://doi.org/10.1093/nar/25.5.0955>.
 12. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glöckner FO. 2007. SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res* 35:7188–7196. <https://doi.org/10.1093/nar/gkm864>.
 13. Nawrocki EP, Eddy SR. 2013. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics* 29:2933–2935. <https://doi.org/10.1093/bioinformatics/btt509>.
 14. Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Pillay M, Ratner A, Huang J, Woyke T, Huntemann M. 2013. IMG 4 version of the integrated microbial genomes comparative analysis system. *Nucleic Acids Res* 42:D560–D567.
 15. Chen IM, Markowitz VM, Palaniappan K, Szeto E, Chu K, Huang J, Ratner A, Pillay M, Hadjithomas M, Huntemann M, Mikhailova N, Ovchinnikova G, Ivanova NN, Kyrpides NC. 2016. Supporting community annotation and user collaboration in the integrated microbial genomes (IMG) system. *BMC Genomics* 17:307. <https://doi.org/10.1186/s12864-016-2629-y>.
 16. Kozłowski JA, Kits KD, Stein LY. 2016. Complete genome sequence of *Nitrosomonas ureae* strain Nm10, an oligotrophic group 6a nitrosomonad. *Genome Announc* 4(2):e00094-16. <https://doi.org/10.1128/genomeA.00094-16>.
 17. Suwa Y, Norton JM, Bollmann A, Klotz MG, Stein LY, Laanbroek HJ, Arp DJ, Goodwin LA, Chertkov O, Held B, Bruce D, Detter JC, Detter JC, Tapia R, Han CS. 2011. Genome sequence of *Nitrosomonas* sp. strain AL212, an ammonia-oxidizing bacterium sensitive to high levels of ammonia. *J Bacteriol* 193:5047–5048. <https://doi.org/10.1128/JB.05521-11>.